

Turf, Lawns, Ornamentals, and Interior Plantscapes

**A Guide for Kentucky Commercial Applicators
in Category 3**



*Kentucky Pesticide Education Program copyright © 2023
University of Kentucky Department of Entomology*

Laws and Regulations

Federal Authority

Pesticides provide important benefits when used correctly. However, they can cause serious harm if used improperly. The **Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA)** is the most important law regulating the registration, distribution, sale, and use of pesticides in the US. However, the Endangered Species Act and Migratory Bird Treaty Act also impact pesticide regulation, application and pest management. FIFRA gives the Environmental Protection Agency (EPA) the authority to oversee the sale and use of pesticides. Commercial applicators can be fined as much as \$5,000 for FIFRA violations. Criminal penalties can be as much as \$25,000 and/or 1 year in prison. In addition, Kentucky can enact legal requirements that may be more restrictive than federal law. In Kentucky, the Kentucky Department of Agriculture administers the EPA-approved certification program and enforces FIFRA regulations.

FIFRA give EPA Authority to:

- Impose civil and/or criminal penalties on anyone who misuses a pesticide or commits any other listed unlawful acts. Fines can be up to \$1,000 for each offense. However, you have the right to ask for a hearing in your own city or county.
- Stop the sale or use of any pesticide.
- Issue removal orders and seize products to keep them out of the market if it determines the products pose an unreasonable risk.
- Reevaluate older pesticides to ensure that they meet more recent safety standards.
- Protect agricultural workers and pesticide handlers from occupational pesticide exposure.

Exceptions to Pesticide Labeling

Unless the label specifically prohibits it, you can apply a pesticide

- To control a pest that is not on the label as long as the specific crop or site is listed
- By any method that is not prohibited. For example, some pesticides cannot be applied by air.
- At a lower dosage, concentration, or less frequently than specified on the label
- In a pesticide-fertilizer mixture.

All pesticides are classified according to their potential hazards under the circumstances in which they are to be used. The two main classifications are **Restricted Use (RUP)** and **unclassified or general use**. The EPA has officially classified very few pesticides as general use. Most that might be expected to fit into the general-use category currently are unclassified. Normally, general-use pesticides have a lower toxicity than RUPs so they are less likely to harm humans or the environment. The general public can buy general-use pesticides without special permits or restrictions. Restricted Use Pesticides can only be purchased and used by a certified applicator in Kentucky.

Endangered Species Act

Plants and animals classified as endangered or threatened must be protected, and this includes from the effects of pesticides. Some pesticides may have specific restrictions on their use in areas of endangered species habitat. This may include special label instructions to check with an EPA website to determine if there are specific precautions to take when using the product in their area.

Kentucky Laws and Regulations

The Division of Environmental Services of the Kentucky Department of Agriculture (KDA) regulates federal and state pesticide laws and regulations, including the Kentucky Fertilizer and Pesticides Storage, Pesticide Use and Application Act of 1996 (KRS 217b) which was revised in 2022. The Kentucky Department of Agriculture implements the provisions of KRS 217b through the administrative regulations, 302 KAR 026. It is responsible for regulating the registration, sale, distribution, proper use, storage, disposal, and application of pesticides in the Commonwealth. The Division strives to educate the pest control industry and consumers about the proper use of pesticides through education and training programs.

KDA personnel give exams to certify and license qualified citizens who wish to apply or to sell pesticides. Field inspectors from the Agricultural Branch inspect facilities of the businesses which sell and/or apply pesticides and review their records. They can impose fines on businesses and/or individuals who neglect to follow federal and state laws concerning the proper storage, containment, sale,

distribution, application, record keeping, or disposal of federally registered pesticides. They also investigate potential pesticide application complaints and violations.

You are responsible for learning about and complying with pesticide laws and regulations before making any applications. In addition, you are responsible for any consequences of actions that result from an application. **Ignorance of the law is never an excuse for noncompliance or violations.**

Important Definitions

- **Application** - placing of a pesticide or pesticide impregnated fertilizers for effect, including mixing and loading.
- **Applicator for hire** - any person who makes an application of lawn chemicals to a lawn for compensation, including applications made by an employee to lawns owned, occupied or managed by his/her employer.
- **Certification** - recognition by the KDA that a person has demonstrated a minimum level of competence by examination and continuing education units and is authorized to use or supervise the use of pesticides in the area of certification.
- **Commercial Pesticide Applicator** - any individual employed by an operator to apply pesticides. Applicators must be certified in the appropriate category and must have a valid license issued by the KDA. The annual applicator license expires on December 31, the license fee is \$25.
- **Commercial Pesticide Operator** - owns or manages a business that applies pesticides on the lands of another for hire. Operators must be certified in the appropriate category and must have a valid license issued by the KDA. A licensed commercial pesticide operator also must be registered as a pesticide dealer or must be employed by a registered dealer. The annual operator license expires on December 31, the license fee is \$100.
- **Customer** - a person who makes a contract, either written or verbal, with an applicator for hire to make an application.
- **Dealer** - stores bulk fertilizer or a restricted use pesticide for redistribution or direct resale, OR is in the business (for compensation) of applying any pesticide to the lands of another.
- **Direct on-the-job supervision** - when a licensed operator or applicator is physically on site and is

directly supervising or training an individual to apply a pesticide.

- **Golf course** - land, including a lawn, on which an application is made for the purpose of maintaining that land for use in the game of golf.
- **Lawn** - land covered with turf, including ornamental plants, maintained for the purpose of human use and enjoyment of outdoor areas.
- **Lawn chemicals** - fertilizers, pesticides, or defoliants applied or intended for application to lawns.
- **License renewal** - There is a 25% fine for license holders who do not file a renewal before January 31. The licensee must take a new certification examination if the license is not renewed before this deadline.
- **Noncommercial applicator** - an employee of a golf course, municipal corporation, public utility, or other governmental agency certified and licensed to apply pesticides to lands owned, occupied, or managed by his or her employer. The annual non-commercial applicator license expires on December 31, the license fee is \$10.
- **Plant-regulating materials** - fertilizers, pesticides, or defoliants applied or intended for application to a golf course.
- **Pests** - any animals (insects, snails, slugs, rodents, etc.); plant pathogens (nematodes, fungi, viruses, bacteria, or other microorganisms) or plants normally considered to be a pest, or which are declared to be a pest by the KDA.
- **Pesticide** - any substance or mixture of substances intended to:
 - prevent, destroy, control, repel, attract, or mitigate any pest;
 - be used a plant regulator, or a spray adjuvant, after being mixed with an EPA registered product;
 - be used as a plant regulator, defoliant, or desiccant.
- **Restricted Use Pesticide** -any pesticide classified as such by the EPA administrator, or by administrative regulation of the KDA. Only certified applicators can purchase and use them.

Generally, the EPA classifies a pesticide as restricted use if:

- it exceeds one or more human health toxicity criteria;
- it meets certain criteria for hazards to non-target organisms or ecosystems, or the EPA

determines that a product (or class of products) may cause unreasonable harm to human health and/or the environment without such restriction;

- then the restricted-use classification designation must appear prominently on the top of the front panel of the pesticide label.
- **Structural pest** - a pest which commonly invades or attacks dwellings or structures.
- **Trainee** - an individual employed by a dealer and working under the direct on-the-job supervision of a licensed operator or applicator. Trainees must be registered with the Kentucky Department of Agriculture with the registration valid for 90 days and cannot be renewed. The fee for trainee registration is \$25.
- **Turf** - the upper layer of soils bound by grass and plant roots into a thick mat.

Recordkeeping Requirements

State law requires that any certified applicator keep records of all applications of general and restricted use pesticides. **Keep the records for at least 3 years** from the date of application. USDA and/or KDA representatives have legal access to the records. Pesticide application records must be recorded within 14 days from the date of application. These records must include:

- name and address of person receiving application services;
- location of application;
- size of area treated;
- crop, commodity, stored product, or type of area treated;
- time and date of application;
- brand name or product name of pesticides applied;
- EPA registration number;
- total amount of each pesticide applied per location per application;
- name of person making the pesticide application;
- if application is made by a trainee, the name of the trainee;
- if application is made by a trainee, name and license number of the supervising applicator;
- records required related to trainee supervision;
- purpose of application; and
- any other record as required by the label.

Pesticide applications records:

- are invaluable documentation in the event of a complaint or lawsuit.

- can help determine which pesticide treatments work, which do not work, and why
- help you to plan purchases so that you buy only the amount needed
- provide information needed by medical staff
- document the steps taken to protect farmworkers and the environment
- are used for federal and state surveys

Posting and Public Notification Requirements

Lawn Care: Immediately following an application to a lawn, place a lawn marker at a prominent location in the lawn. The lawn marker must consist of, at a minimum, a four (4) inch by five (5) inch white sign attached to the upper portion of a dowel or other supporting device of not less than twelve (12) inches in length. Lettering on the lawn marker must be in a contrasting color and must read on one side "LAWN CARE APPLICATION - PLEASE STAY OFF GRASS UNTIL DRY" in letters easily readable and not less than three-eighths (3/8) inches in height. The lawn marker may also display a symbol depicting the required message and the name, logo, and service mark of the applicator. The lawn marker must be removed and discarded by the property owner or resident, or other person authorized by the property owner or resident, the day following application. **For applications to residential properties of three (3) families or less, the applicator must place one (1) lawn marker per property. For applications to properties other than residential property of three (3) families or less, the applicator must place lawn markers at primary points of entry to the property** to provide notice that lawn chemicals have been applied to the lawn.

Ornamental Care: Immediately following an application of pesticides to ornamentals, place a marker at a prominent location in the ornamentals that reads "PESTICIDES HAVE BEEN APPLIED - PLEASE STAY OUT OF TREATED AREA" in letters easily readable and not less than three-eighths (3/8) inches in height. The marker may also display a symbol depicting the required message and the name, logo, and service mark of the applicator.

Provide prior notification to the customer or adjoining residents in writing, in person, or by telephone. If requested, of the date and approximate time of the application. If you are not able to provide prior notification to a customer or adjoining residence due to the absence or inaccessibility of the individual, leave a written notice at the residence.

Golf Courses: Immediately following the application of plant-regulating materials on a golf course, the applicator must place a golf course marker on the number-one (1) and number-ten (10) tees. The **golf course marker** consists of, at

a minimum, a four (4) inch by five (5) inch white sign attached to the upper portion of a dowel or other supporting device of not less than twelve (12) inches in length. Lettering on the golf course marker must be in a contrasting color and must read on one side "PLANT-REGULATING MATERIALS HAVE BEEN APPLIED. IF DESIRED, YOU MAY CONTACT THE GOLF COURSE SUPERINTENDENT FOR FURTHER INFORMATION" in letters easily readable and not less than three-eighths (3/8) inches in height. The golf course marker may also display a symbol depicting the required message and the name, logo, and service mark of the applicator. The golf course marker may be removed by the applicator or other personnel authorized by the golf course management the day following application.

Any person whose residence directly adjoins a golf course may request prior notification of an application by contacting the golf course superintendent's office and providing his or her name, address, and telephone number. If requested, the golf course must provide notification in writing, in person, or by telephone. In the event the golf course cannot provide advance notice, the person must be contacted at the time of application. If the golf course is unable to provide prior notification or direct notification to a resident because of the absence or unavailability of the resident, the golf course must leave a written notice at the residence.

Sports Turf: Immediately following an application to turf on a sports field, the applicator must place a marker at usual entry points to the field. The marker must consist of, at a minimum, a 4 in. x 5 in. white sign attached to the upper portion of a dowel or other supporting device of not less than twelve (12) inches in length. Lettering on the marker must be in a contrasting color and must read on one (1) side "PESTICIDES HAVE BEEN APPLIED - PLEASE STAY OUT OF TREATED AREA" in letters easily readable and not less than three-eighths (3/8) inches in height. The marker may also display a symbol depicting the required message and the name, logo, and service mark of the applicator. The marker may be removed by the applicator or other personnel authorized by the sports field management the day following application.

Any person whose residence directly adjoins a sports field may request prior notification of an application by contacting the sports field manager's office and providing his or her name, address, and telephone number. If requested, the manager must provide notification in writing, in person, or by telephone. In the event the sports field manager cannot provide advance notice, the person must be contacted at the time of application. If the manager is unable to provide prior notification or direct

notification to a resident because of the absence or unavailability of the resident, the manager must leave a written notice at the residence.

Interior Plantscapes: Immediately following an application to interior plantscapes, place a marker at a prominent location in the interior plantscapes. The sign must read "PESTICIDES HAVE BEEN APPLIED - PLEASE STAY OUT OF TREATED AREA" in letters easily readable and not less than three-eighths (3/8) inches in height. The marker may also display a symbol depicting the required message and the name, logo, and service mark of the applicator. Posting requirements must not apply if plants that are in interior plantscapes are taken off-site for an application and not returned until the plants have adequately dried.

Provide prior notification to the customer or adjoining residents in writing, in person, or by telephone if requested, of the date and approximate time of the application. If an operator is not able to provide prior notification to a customer or adjoining residence due to the absence or inaccessibility of the individual, the applicator must leave a written notice at the residence.

Customer Notification Requirements

Any operator or applicator making lawn care applications must provide the customer at the time of entering into a contract and at the time of application to a lawn with written information concerning lawn chemicals, application procedures, and other general guidelines. An applicator shall provide the following information upon request to all persons requesting notification, and must record and maintain at the applicator's business address the following information relating to the application of each pesticide used:

The application records listed above must be maintained in the golf course superintendent's or the sports turf manager's office and be readily available to review on request. This record shall be retained for at least three (3) years and be available for inspection by the Kentucky Department of Agriculture.

- brand name or common name of the pesticide applied;
- pesticide type;
- fertilizer rate and analysis;
- reason for use;
- concentration of end use product applied;
- rate of application,
- total gallons of end use product applied;
- any special instruction appearing on the label of the pesticide product applicable to the use of the treated area following application;
- any other precautionary or hazard information appearing on the label as applicable to the end use concentration;
- name and applicator license or certificate number of person actually making the application;
- customer name, address, and date of application;
- location area of area treated; and
- total area treated.

Certification and Licensing

Commercial and non-commercial pesticide applicators must be both certified and licensed. Both are accomplished by passing a written test (minimum score 70%) administered by the KDA. The test is based on information in this manual.

Evidence of Financial Liability

Pesticide dealers who apply pesticides to the lands of others must show evidence of financial responsibility. This can be a surety bond or a liability insurance policy of at least one million dollars (\$1,000,000) that would protect persons who may suffer legal damages as a result of the applicant.

How To Remain Licensed and Certified

1. Return the annual license renewal form before January 31. There is a 25% penalty added to the original fee for license holders who do not renew and pay their fees before January 31. Failure to renew a license by January 31 of each year, will also result in the former license holder being required to retest as an initial applicant, after any applicable fines are paid.

2. Pay any required fees.
3. Earn Continuing Education Units (CEUs) in educational meetings approved by the KDA. Twelve CEU credits, with at least one related to each category of license held by the person within the three-year period prior to each annual license renewal application.

The Kentucky Cooperative Extension Service provides training materials and educational programs for certification and continuing education of commercial and non-commercial applicators through the Pesticide Safety Education Program.

Penalties

Anyone who uses a pesticide in a manner inconsistent with its labeling directions and restrictions may be subject to civil and/or criminal penalties. Generally, any applicator in violation of FIFRA may be assessed a civil penalty. However, the EPA may issue a warning instead of assessing a penalty. An intentional violation by a private applicator is a misdemeanor and will result in a fine and/or up to 30 days imprisonment. You must use all pesticides exactly according to labeling directions—the label is the law!

Practice Questions

1) The _____ is the most important law regulating pesticides in the US.

1. KRS 217b Ky Fertilizer and Pesticides Storage, Pesticide Use and Application Act of 1996
2. 1996 Farm Bill
3. Federal Insecticide, Fungicide, and Rodenticide act (FIFRA)
4. Ky Department of Ag Regulation 1262

2) Commercial and non-commercial pesticide operator and applicator licenses are good for ____ year(s).

1. 1
2. 3
3. 5
4. 10

3) Commercial and non-commercial pesticide applicator certifications are good for ____ year(s).

1. 1
2. 3
3. 5
4. 10

4) A pesticide is categorized as general use if it can harm humans or the environment even if it is used according to label directions.

1. True
2. False

5) A minimum score of ____ % is required on the test to become a commercial or non-commercial pesticide applicators.

1. 60
2. 70
3. 80
4. 100

6) According to state laws and regulations, anyone who is in the business of applying any pesticide to the lands of another is considered to be a pesticide dealer.

1. True
2. False

7) _____ applicators are people who apply pesticides to lands owned, occupied, or managed by a golf course, municipal corporation, public utility, or other governmental agency.

1. Certified commercial
2. Registered
3. Non-commercial

8) Non-commercial applicators may apply pesticides to residential or commercial properties for hire without any additional certification.

1. True
2. False

9) A certified commercial or non-commercial pesticide applicator can stay certified by earning _____ continuing education units (CEUs) before their certification expires.

1. 12 CEU hours with at least one in the category held
2. 9 general and 3 specific CEU hours
3. 12 CEU hours in each category held
4. none, you must take a test every 3 years

10) According to Kentucky pesticide laws and regulations, commercial and non-commercial applicators must keep records of both general and restricted use pesticide applications.

1. True
2. False

11) A certified pesticide operator or applicator who fails to renew his/her license before _____ must take a new examination.

1. January 31
2. March 1
3. June 1
4. November 30

12) Fertilizers are not considered to be lawn chemicals.

1. True
2. False

13) Fertilizers, pesticides, or defoliants applied or intended for application to a golf course are defined as _____.

1. general use pesticides
2. restricted use pesticides
3. plant-regulating materials

14) An applicator for hire shall provide the customer with information about lawn chemicals used _____ of application.

1. 24 hours before
2. within 14 days
3. within 30 days
4. at the time of

15) If requested, an applicator for hire shall notify a neighbor whose residence adjoins that of a customer _____ an application of lawn chemicals.

1. 24 to 48 hours before
2. 24 to 48 hours after
3. 7 days before
4. at the time of

16) The applicator shall place a lawn marker at a prominent location in the lawn _____ an application.

1. 24 to 48 hours before
2. immediately after
3. 7 days before
4. 24 to 48 hours after

17) The minimum size of a lawn marker in inches is ____.

1. 2 X 3
2. 3 X 4
3. 4 X 5
4. 5 X 6

18) The minimum size of a golf course marker in inches is ____.

1. 2 X 3
2. 3 X 4
3. 4 X 5
4. 5 X 6

19) _____ lawn marker(s) must be placed on each residential properties of three (3) families or less.

1. One
2. Three
3. None are needed

20) A lawn maker must remain in place for 7 days following an application.

1. True
2. False

21) The golf course marker may be removed by the applicator or other personnel authorized by the golf course management the day following application.

1. True
2. False

22) Immediately following the application of plant-regulating materials on a golf course, the applicator shall place a golf course marker on the _____.

1. 1 and 10 greens
2. 1 and 10 tees
3. 1 and 10 roughs
4. clubhouse lawn

Answers

1: 3 2: 1 3: 2

4: 2 5: 2 6: 1

7: 3 8: 2 9: 1

10: 1 11: 4 12: 2

13: 3 14: 4 15: 1

16: 2 17: 3 18: 3

19: 1 20: 2 21: 1

22: 1

Weeds of Ornamentals and Turfgrass

Weeds are plants that are growing where they are not wanted. Usually, they cause minimal problems in healthy, vigorously growing turf. However, weeds can out-compete desirable turf due to poor management practices: irrigation, fertilization, mowing, turf diseases, insects, or from heavy use. Weed free turf is not practical in most cases but a balanced management program can keep them to a minimum.

A typical weed has one or more of the following characteristics:

- Produces lots of seed
- Populations establish rapidly
- Seeds can lie dormant for a long time
- Have vegetative reproductive structures
- Adapted for easy spread
- Plant development stages

Plant Development Stages

Most plants undergo four stages of growth and development.

1. **Seedlings** emerge from the soil soon after germination.
2. Leaves, stems, and roots grow rapidly during the **vegetative** stage, water and nutrient demands are relatively high.
3. After a period of vegetative growth, the plant enters the **reproductive** stage where most of the energy production in the plant is devoted to seed formation. Seed production is critical for survival of annual and biennial species.
4. Little or no energy production occurs during **maturity** when seed production is nearly finished. During this stage, the plant typically sheds its seeds and dies.

Plant Life Cycles

Annual plants complete their life cycle in one growing season, often in as little as 45 days. **Biennials** require two seasons. **Perennials** grow for three or more years.

Annuals that grow from **spring to fall** (large crabgrass and goosegrass) are **summer or warm-season annuals**. These are often problems because their life cycle is the same as many crops. Those that grow from **fall to spring** (common

chickweed and henbit) are **winter or cool-season annuals**.

Biennial plants complete their life cycles over **two growing seasons**. Most start from seed in the fall or spring and grow through the summer, fall, winter, and following spring. They overwinter as rosettes. In the second summer, biennials flower and die. Examples include wild carrot and musk thistle.

Perennials often are the most difficult weeds to manage. Woody species generally go dormant in the winter and begin growth in spring from aboveground stems. Aboveground parts of herbaceous perennials may die back, but their underground storage organs survive the winter. Many are deep rooted so they continue to grow during summer droughts. Perennials can spread from seed and often from roots, tubers, bulbs, and rhizomes. White clover and yellow nutsedge are examples. Dandelions can be annual or perennial.

Many weeds produce large quantities of seeds that are easily carried by wind, rain, machinery, animals, and people. Weed seeds can germinate after being dormant for long periods of time. They also can tolerate extremes in weather such as temperature and moisture. **It is best to control weeds before they produce seeds.**

Plant Classification

Weeds can be grouped into the following categories:

- grasses
- sedges
- lilies
- broadleaves.




Grasses have only one leaf as they emerge from the soil. Their leaves are two-ranked and typically upright, narrow with parallel veins. Grass stems are round and hollow. The root system of a grass is fibrous with the growing point located at or below the soil surface (surrounded by several layers of leaves). Perennial grasses can produce

new shoots from growing points located on rhizomes (belowground) and/or stolons (aboveground).

Sedges resemble grasses but they have triangular stems with three rows (ranks) of leaves. Typically, sedges are listed under the grass section of an herbicide label. They prefer moist, poorly drained soils, but can grow in fertile, well-drained soils. Yellow nutsedge, is a perennial that reproduces by tubers and rhizomes. It is the principle sedge found in Kentucky.

Lilies resemble grasses and sedges but they have long, linear leaves and reproduce from underground bulbs. Two common species found in Kentucky are wild garlic and Star-of-Bethlehem.

Broadleaf seedlings have two leaves (cotyledons) as they emerge from the soil. The leaves are generally broad with net-like veins. Broadleaves typically have a taproot surrounded by a relative coarse root system. Actively growing broadleaf plants have exposed growing points at the end of each stem and in each leaf axil. Perennial broadleaves may have growing points on roots and stems above and below the surface of the soil.



Summer Annuals

Crabgrass

From seed and roots at lower joints; encouraged by alternating wet, dry soil surface in spring; germination begins mid-April; 3 to 10 finger-like branches at top of stem; thrives in sparse turfgrass stand; low mowing; heavy traffic area; full sun



photo: Lynn Sosnoskie, Univ. of Georgia, Bugwood

Foxtail

Bright green clumping grass; heavy traffic areas; bushy, cylindrical seed head at top of stem; full sun; low mowing



photo: John D. Byrd, Mississippi State Univ., Bugwood.org

Goosegrass

Germinates May to June; prostrate growth; white center with wagon-wheel appearance; sparse turfgrass stand; low mowing; grows well in compacted soil



photo: Joseph M. DiTomaso, Univ. of California - Davis, Bugwood.org

Common knotweed

Germinates in early spring; prostrate growth; alternate, oblong, pointed leaves; heavy traffic; along roads or driveways



photo: Lynn Sosnoskie, Univ. of Georgia, Bugwood.org

Winter Annuals

Annual bluegrass

Germinates from fall to early spring; light green bunch-type grass with flattened stems; can produce seed heads even with low mowing heights; irrigated turf and moist shade



photo: Joseph M. DiTomaso, University of California - Davis, Bugwood.org

Common chickweed

From seed in autumn and creeping stems; pairs of smooth, egg-shaped leaves; small star-shaped white flowers



photo: John D. Byrd, Mississippi State Univ., Bugwood.org

Henbit

From seed and roots form lower joints; square stem that branches close to ground; almost circular opposite leaves with rounded teeth or lobes; prefers moist soil in shade



photo: Chris Evans, University of Illinois, Bugwood.org

Perennials

Broadleaved Plantain

From seeds and new shoots from taproot; leafless stem; egg-shaped leaves; seeds along half length of seed stalk; thrive in weak, thin turf



photo: Chris Evans, University of Illinois, Bugwood.org

White clover

From seed and creeping stems; creeping growth; white blossoms; leaves with 3 leaflets; survives close mowing



photo: Chris Evans, University of Illinois, Bugwood.org

Wild garlic

From underground bulbs and aboveground bulblets; round slender leaves are hollow; poorly maintained or thin turf



photo: Ohio State Weed Lab, The Ohio State University,
Bugwood.org

Yellow nutsedge

From seed, rhizomes, and tubers; fast growing; yellow-green triangular stem; often in wet soil



photo: Howard F. Schwartz, Colorado State Univ., Bugwood.org

Nimblewill

Warm season perennial, stolons root at nodes producing dense stands. Leaf blades gray-green with loose spreading growth. Especially common in Kentucky bluegrass. Mowing is not an effective control measure.



photo: msurfweeds.net

Perennial or Annual

Common dandelion

From seeds and root shoots; stems contain a milky juice and arise from a long, thick, fleshy taproot



Photo: Howard F. Schwartz, Colorado State University,
Bugwood.org

Wild violet

From seed and underground root; heart-shaped leaves; often grow in shade



photo: Rob Routledge, Sault College, Bugwood.org

Practice Questions

1) _____ plants germinate in the spring, grow, mature, produce seed, and die in summer or early fall.

1. Summer annual
2. Sedge
3. Winter annual
4. Broadleaf

2) _____ plants appear in the fall and die in late spring or early summer of the following year.

1. Summer annual
2. Sedge
3. Winter annual
4. Broadleaf

3) Annual plants are easiest to control in the _____ stage.

1. seedling
2. vegetative
3. reproductive
4. mature

4) The _____ stage of plant development is characterized by rapid uptake of water and nutrients and fast growth.

1. seedling
2. vegetative
3. reproductive
4. mature

5) Biennial plants are easiest to control in the _____ stage.

1. seedling
2. vegetative
3. reproductive
4. mature

6) Removing the aboveground vegetation will stop perennial plants from spreading in a field.

1. True
2. False

7) _____ plants can reproduce by stolons, rhizomes, tubers, or bulbs.

1. Annual
2. Biennial
3. Perennial

8) The rosette stage is part of the live cycle of a _____.

1. biennial weed
2. pathogenic fungus
3. sap-feeding insect
4. winter annual weed

9) Which of the following is not a form of vegetative reproduction of a weed?

1. seed
2. stolon
3. rhizome
4. bulb

10) Crabgrass _____.

1. is a winter annual with a bushy seed head
2. is a summer annual with 3 to 10 finger-like branches at the top of its stem
3. is a perennial plant
4. has a square stem

11) Common knotweed _____.

1. grows in low-lying wet areas
2. grows in areas compacted by heavy traffic
3. is a perennial plant
4. has a bushy, cylindrical seed head at the top of stem

12) Annual bluegrass _____.

1. is a light green bunch-type grass that can produce seed heads even with low mowing heights
2. has egg-shaped leaves and small star-shaped white flowers
3. has a prostrate growth form and a white center with a wagon-wheel appearance

13) _____ has a yellow-green triangular stem and grows in wet soil.

1. prostrate knotweed
2. wild garlic
3. yellow nutsedge
4. wild violet

14) An emerging _____ plant has 2 seed leaves.

1. grass
2. broadleaf
3. sedge

15) Annual plants reproduce from underground stems and tubers.

1. True
2. False

16) It is best to control weeds before they produce seeds.

1. True
2. False

17) The growing point of a grass is always above the soil surface.

1. True
2. False

18) All actively growing _____ have exposed growing points at the end of each stem and in each leaf axil.

1. sedges
2. broadleaves
3. grasses

19) _____ usually have a taproot and a relatively coarse root system.

1. Broadleaves
2. Grasses
3. Sedges

Answers

- 1: 1
- 2: 3
- 3: 1
- 4: 2
- 5: 1
- 6: 2
- 7: 3
- 8: 1
- 9: 1
- 10: 2
- 11: 2
- 12: 1
- 13: 3
- 14: 2
- 15: 2
- 16: 1
- 17: 2
- 18: 2
- 19: 1


Pests of Ornamentals and Turfgrass

Insects and Other Arthropods


Insects, spiders, scorpions, millipedes, centipedes, ticks, and mites are arthropods. They have hard external skeletons and segmented legs and bodies. Most insects have 3 main body regions and 3 pairs of legs; they are the only arthropods that can fly.

Insect Life Cycles and Growth

Insects go through a series of changes during their development from egg to adult in a process called **metamorphosis**. When the insect hatches from an egg, it is either a **nymph** (gradual metamorphosis) or a **larva** (complete metamorphosis). The immature stage must shed its external skeleton, a process called **molting**, in order to grow.



Grasshoppers undergo **gradual metamorphosis**, passing through three stages of development: egg, nymph, and adult. Nymphs resemble adults. They eat the same food and live in the same environment. The change in form from nymph to adult is gradual. Only the adult state has wings. Other examples are aphids, stink bugs, and leafhoppers.



Insects with **complete metamorphosis** include butterflies and moths, beetles, flies, bees, and ants. There are **four stages in complete metamorphosis** – **egg, larva, pupa, and adult**. The larvae, are specialized for feeding and look very different from the adult. They have general names such as caterpillar, maggot, white

grub, or wireworm. **Larvae usually live in very different situations and often feed on different foods than adults.**

A variety of insects and mites can attack plants but most are not pests. Some are beneficial, providing natural control or pollination services. Others are scavengers on dead or dying plants so they recycle nutrients. **Just because an insect is around damage does not mean it was the cause.**

Mouthparts and Feeding - Ways Insects Can Damage Plants

Pest insects may be divided into major groups according to how they feed:

1. piercing-sucking
2. chewing
3. rasping plant tissue

Sap feeders with piercing-sucking mouthparts can cause wilting, leaf curl, or stunted foliage. Chemicals injected by some species of leafhoppers can cause leaf burn. Stink bug feeding can cause distorted leaves or fruit. Several aphid and leafhopper species can carry virus diseases.

Rasping tiny thrips tear plant cells and feed on sap. These tiny insects may leave feeding scars or distorted leaves; some can carry plant disease.

Chewers include caterpillars and beetles. They feed on foliage. The amount of feeding a plant can tolerate without significant impact on growth or yield varies with a plant's age, growth stage, or stress (drought, etc.).

Arthropod Pests of Trees and Shrubs

Sap Feeders (Piercing-Sucking)

Pests with sucking mouthparts cause similar types of damage. Using their mouthparts, the pests pierce or rasp tissue so they can suck plant juices. **Damaged foliage is usually mottled, but other symptoms may be wilting, scorched leaf tips, or puckering and curling.** When sooty mold occurs on plants, it is almost always associated with "honeydew" that is excreted by certain kinds of sucking insects.



Aphids, photo: UK Entomology

Aphids or “plant lice,” are small, soft-bodied insects that usually cluster on stems or undersides of terminal leaves. Aphids may be green, black, or red, but sometimes their color is hidden by a white waxy coating. Much of the sap that aphids suck passes through them undigested and is excreted as “honeydew.” Honeydew makes the leaves sticky, and sooty mold may grow on these deposits. Feeding by some kinds of aphids will cause leaves to pucker, curl, or twist.



Calico scale, photo. UK Entomology

During most of their lives, **scale insects** are legless and motionless and do not resemble insects at all. They may be circular, oval, or pear-shaped. Some are flat, others convex. **Two major groups of scales are most common in Kentucky.** The **armored scale** produces a waxy shell that gives the soft-bodied insect under it some protection. The **soft scales** do not produce a shell, but their bodies may be tough.

Scales reproduce by giving birth to “crawlers” or by laying eggs that hatch into crawlers. **Crawlers** have legs, eyes, and antennae, all of which allow them to move out from under the mother’s shell or body and seek a suitable place of their own on the plant. Soon after inserting their beak to feed, they molt and lose their legs, eyes, and antennae and remain motionless for the rest of their lives.

Plants infested with scales may lack vigor and appear sickly. Soft scales, like aphids, produce honeydew and cause the same symptoms as mentioned for aphid honeydew. **It is**

easier to control most scales while they are in the crawler stage because they are not protected by a shell or waxy coat. Treatment applied for scale control should coincide with crawler activity. A second treatment in 2 to 3 weeks often is recommended. Timing of systemic insecticide applications is not so critical.

On oil-tolerant plants, **oil sprays** can be used to control all stages of scales, including eggs. Summer oils may be effective during the warmer months. Apply dormant oils in winter. **Insecticidal soaps** are another alternative for controlling scale crawlers as well as aphids, mealybugs, whiteflies, thrips, and mites.



Spider mite, photo: Washington State University

Mites are not insects, but their damage and the methods of control are similar to those of insects. They differ from insects in that they have 8 legs, not 6, and have only 1 body region instead of 3. All mites are tiny and usually cannot be seen without the aid of a magnifying lens. **By tapping infested twigs over a sheet of white paper, the dislodged mites are much easier to detect.** They vary widely in color. Some mites spin fine, delicate webbing on the host plant. This webbing is usually easier to detect than the mites themselves.

Spider mites feed by sucking cell contents from individual leaf cells. Initial damage appears as fine light dots on the leaves. A small number of mites usually is not reason for concern but very high populations can cause significant damage.

Mite **damage often appears as a bronzing of the foliage**, which sometimes gives it a dusty appearance. Leaf drop may also occur. As feeding continues, the leaves turn yellowish or reddish and drop off. Often, large amounts of webbing cover infested leaves. Damage is usually worse when plants are under drought stress.

Many kinds of plants are attacked by the **two-spotted spider mite**, and almost all coniferous plants are hosts to the **spruce spider mite**. The **Southern red mite** is primarily a pest on broadleaf evergreens such as azaleas and camellias.

Foliage Feeders (Chewing mouthparts)

Beetles vary considerably in size, shape, color, and habits but all have chewing mouthparts. One of their most distinctive features is that their front wings are hard or leathery and meet in a straight line down the center of the back.

Beetles may attack any part of a plant and in various ways. Some are typical leaf feeders and bite off pieces of leaf, while others are leaf miners or skeletonizers. With some beetles, the adults and larvae both are leaf feeders on the same plant; other beetles may be foliage feeders as adults and root feeders on other plants while in the larval stage. Some feed during the day and some feed only at night, such as the May beetles.

Beetles



Japanese beetle, photo: UK Entomology

The adult **Japanese beetle** causes serious damage to the foliage of many landscape plants. The larvae stage of the Japanese beetle is a **white grub** that feeds below ground on plant roots and is a serious pest of turfgrasses.

Some beetles, such as the **bronze birch borer**, feed as the larval stage in the cambium of trees and shrubs. This boring activity leaves “galleries” underneath the bark, usually causing serious damage to host plants. Girdled plants usually die.

Caterpillars

Caterpillars are the worm-like immature stages of moths and butterflies. They range in size from tiny to 5 inches long. They usually have a distinct head and 4 pairs of fleshy legs on the middle of the body. The body may be fuzzy, naked and smooth, or spiny.

Caterpillars are primarily foliage feeders and eat out irregular areas or they may entirely strip the leaves. Some caterpillars, because of their special habits, are also referred to as webworms, tent caterpillars, leaf rollers, leaf folders,

skeletonizers, bagworms, and leafminers. Some feed as individuals; others feed in groups or colonies.



Fall webworm, photo: Steven Katovich, USDA Forest Service, Bugwood.org

Fall webworms build tents at ends of branches. When only a few large caterpillars are present, handpicking is effective. **Webworms and tent caterpillars can either be pruned out or burned out with a torch.** If pruning would adversely affect a plant or if the infestation of any caterpillar is generally distributed over a plant, **a single treatment of an approved insecticide applied when the caterpillars are young** will usually give control.



Clearwing borer moth, photo: UK Entomology

A group of small moths, usually referred to as **clearwing moths**, cause serious boring damage to certain plants. The active adults often resemble wasps. The larvae bore through the cambial layer, causing stress, decline and, occasionally, death of plants. **Dogwoods, lilacs, and ash are affected by clearwing borers.**

Sawflies are wasp-like insects and are related to typical wasps, bees, and ants. The larval stages of most sawflies resemble naked caterpillars, but they have more than 5 pairs of fleshy legs on the body while caterpillars have only 4 or fewer pairs. Some sawfly larvae are slug-like in appearance, such as the pear slug and rose slug.



Slug-like sawfly larvae, photo: UK Entomology



Redheaded pine sawfly, photo: UK Entomology

Most sawfly larvae are foliage feeders that eat the entire leaf but **slug sawflies** are skeletonizers. A few types of sawflies are wood borers or leafminers. These differ further from typical sawflies in that they do not have fleshy abdominal legs. The **most serious sawfly pests in our area are those that attack coniferous shrubs and trees**. They feed in groups and can quickly defoliate a plant. This defoliation often leads to the plant's death. A single application of an approved insecticide is usually sufficient for control.

Borers

Wood-boring insects are among the most destructive pests of ornamental trees and shrubs. Borers are the larvae, or immature stage, of certain moths and beetles. They tunnel and feed under the bark in living wood, destroying water- and sap-conducting tissues. This causes girdling, branch dieback, structural weakness, and decline and eventual death of susceptible plants. Infestation sites also provide entry points for plant pathogens.

Clearwing (moth) and flatheaded borers (beetles) are the two main types that attack woody ornamentals.




Flatheaded borer and tunnels, photo: Eric R. Day, Virginia Polytechnic Institute and State University, Bugwood.org



Flatheaded borer, photo: UK Entomology

Flatheaded borers are so named because their first body segment, behind the head, is flattened laterally. The adults are fast-moving, flattened, metallic-colored beetles with short antennae. The whitish, legless larvae make winding tunnels beneath the bark, destroying phloem and cambium and girdling the trunk or branches. The tunnels may be visible externally as spiral ridges or cankers on the limbs or trunks. Unlike clearwing borers, which expel frass from cracks in the bark, **flatheaded borers pack their fine, sawdust-like frass in their tunnels**. Examples are the emerald ash borer, bronze birch borer, and flatheaded appletree borer.



Dogwood borer larva, photo: UK Entomology

Clearwing borer larvae are whitish, hairless caterpillars with a brown head. Adults are delicate, day-flying moths that resemble small wasps. The moths feed only on nectar or not at all, so they do not cause damage. There are a number of different species, but **the most damaging clearwing borers are associated with dogwood, lilac, ash, oak, rhododendron, and ornamental Prunus species, including flowering peach, plums, and cherries.**

These groups differ somewhat in their habits and host preferences, but similar management tactics are used for both. The keys to controlling these pests are to keep plants healthy and, if necessary, to treat during those brief times of the year when the insects are vulnerable to insecticides.

Borers rarely infest healthy plants growing in their natural environments. However, **when trees or shrubs are transplanted into the landscape, stresses such as drought, soil compaction, sun scald, or injuries can weaken them and make them more susceptible to attack.** Research has shown that the adults may locate suitable egg-laying sites by responding to volatile chemicals that emanate from stressed trees. **The invasive emerald ash borer is an exception. It attacks healthy ash trees.**

Adult borers emerge from infested trees in the spring or summer. After mating, the females fly to a suitable host and lay eggs on the bark, often in crevices or around wounds. Hatching occurs about 10 days to 2 weeks later, and the young larvae quickly tunnel beneath the bark where they feed and grow. **Once inside the tree, borer larvae are no longer vulnerable to insecticide sprays and are seldom detected until serious damage has been done.** Systemic insecticides can protect some trees and shrubs from attack by some borers.

Several species of clearwing and flatheaded borers can infest landscape plants. While some are attracted to a wide range of hosts, most attack only particular kinds of trees and shrubs. **In order for treatment to be effective, it is important to know when the adults of each species are active and which plants are vulnerable.**

Gall Makers



Horned oak gall, photo: UK Entomology

When some insects or mites lay eggs in tissue, they may inject a chemical into the plant that causes it to grow abnormally, producing a **gall**. Plant parts affected include roots, crown, bark, branches, twigs, buds, and leaves. **Each species of insect or mite produces a characteristic gall on host plants.** With the exception of horned and gouty oak galls, **most galls do not harm tree health.**

Insect Pests of Turf

Lawns often include a variety of insects, some of which are direct pests of grass, or nuisances and pests to humans and pets. Some may be predators or parasites of other insects, or harmless scavengers. Through complex interactions between the insects and other factors, the lawn ecosystem becomes more or less balanced. If we are not satisfied with the balance, we may use maintenance practices to improve our lawns.

However, the solution may trade one problem for another. For instance, fertilization to increase grass lushness may favor the development of certain insect and disease problems. Insecticidal control for one kind of insect may kill predators or alter competition, allowing a different insect pest to flourish. **Often, the side effects of management practices cannot be precisely predicted, so lawn situations need to be monitored over time and maintenance practices modified, if necessary.** Some of these interactions and problems are demonstrated in the case of white grubs as lawn pests in Kentucky.



White grub, photo: UK Entomology

White grubs are the larval stages of scarab beetles such as masked chafers, rose chafer, May beetles, green June beetle, and Japanese beetle. White grubs look more or less alike. They have brown distinct heads and thoracic legs, and the body is whitish, fat, and usually curled into a C-shape. Size varies from 1/8 to 1-1/2 inches long depending on the age and species. The grubs occur in large patches of sod an inch or so below the soil line where they consume the anchoring roots of grass. During dry weather, the infested sod may die for lack of water.

Soil insecticides for white grub control should be applied in August before the grubs cause serious damage. Most instances of control failures are a result of poor timing or techniques of insecticide applications.



photo: Michigan State University Extension

Sod webworms graze baseball-sized patches of grass that turn brown and die. Patches of grass that are clipped off at the soil surface may be numerous and run together to form large dead areas. **Dirty silk tubes containing the inch-long caterpillar or pupa can usually be found in the thatch of killed spots.** The **adult stage of the pest is a small buff moth** that is often seen fluttering over lawns at dusk and at night around lighted doorways about two weeks before larvae become numerous. There are up to three generations per year.



Fall armyworm, photo: UK Entomology

Armyworms, including the true armyworm and the fall armyworm, are characteristic caterpillars about 1-1/2 inches long when full grown. They vary in intensity from year to year, but during outbreaks they may move across an area in army fashion completely stripping grasses in their path. Fescue is more often attacked than bluegrass. These insects are also important pests of grain crops.



Bronze cutworm, photo: UK Entomology

Various species of cutworms occur in turf and some are hard to distinguish from armyworms based on body characteristics. However, they never occur in large numbers as do armyworms.


Beneficial Insects

Beneficial insects can help to regulate pest populations. Predators generally are not very selective and may feed on non-pest species. Some species of wasps and flies are very selective in the prey that they attack including leaf feeding caterpillars and wood boring beetles. Four species of lady beetles feed on soft-bodied insects such as aphids and scale crawlers.



Pink lady beetle, photo: UK Entomology

The pink spotted lady beetle has a medium-sized, oblong pink to red body marked with black spots. Adults and larvae are important aphid predators but they also eat mites, insect eggs, and small larvae. Unlike most lady beetles, plant pollen may make up to 50% of the diet.



Multicolored Asian lady beetle, photo: UK Entomology

Multicolored Asian lady beetle is a large orange lady beetle with a variable number of spots. It can be recognized by the black 'M' on the white segment over the head. Aggregations of these insects find their way into homes in the fall, where they are a nuisance. Their secretions can ruin rugs and other furniture, but they do not breed or feed inside the home.



Convergent lady beetle, photo: UK Entomology

The convergent lady beetle is an important natural enemy of aphids, scales, thrips, and other soft-bodied insects. It will also feed on pollen and nectar from flowers when prey is scarce. Larger larvae are voracious feeders and may consume between 30 and 50 aphids per day.



Seven-spotted lady beetle, photo: UK Entomology

The seven-spotted lady beetle is a medium-sized, orange beetle with seven black spots. It is a European species that was introduced into the US to aid in managing some aphid pests.



Jumping spider with prey, photo: UK Entomology

Spiders are general predators that feed on many types of prey including pest and beneficial species.



*Beneficial wasp laying its eggs in a caterpillar
Photo: S. Bauer, USDA*

The larvae of many beneficial wasps develop in the bodies of caterpillars. Adult wasps can be very selective when choosing prey.

Other Pests



Worm castings on golf course, photo: Penn State Univ.

Earthworms and nightcrawlers are not insects but they can be abundant in soil. These annelids help to recycle organic matter and their burrows allow oxygen and water to enter the soil more easily. However, when many are present, their activities and casting can cause the surface to be very lumpy. In addition, new species (green stinkworms, Asian yellowworms) are beginning to appear in some locations. These worms can cause significant problems in lawns, athletic fields and on golf greens.

Earthworms and night crawlers are generally found in the top 12" to 18" of the soil because this is where food is most abundant. They swallow soil and organic matter and grind it in the gizzard section of the digestive tract. Undigested material (castings) is used to line the burrow or is eliminated on the surface. They are most active when the soil is warm and moist. The worms move deeper as the soil dries in summer. Control strategies for these pests are being developed.



photo: Alfred Viola, Northeastern University, Bugwood.org

Skunks damage turf when they discover abundant white grub populations. Skunks dig through the sod and feed on the white grubs, thereby uprooting the sod and aggravating the damage already begun by the grubs. Skunks also may

spray a disagreeable smelling substance on unwary people or pets who disturb them.



photo: Terry L. Spivey Photography, Bugwood.org

Birds, especially crows, starlings, and grackles, commonly tear up infested turf in search of grubs. Flocks of blackbirds frequenting a turf site, or holes left in the turf by their beaks, may indicate a grub problem.



Mole, photo: UK Entomology

Moles feed primarily on earthworms, but they may also feed on white grubs, wireworms, beetles, and many other invertebrates. They do not feed on plant roots or other underground plant growth.

However, as they tunnel along in their surface runs, moles damage turf roots and may destroy newly seeded lawns. In established turf, the mower may skin the tops of the runs and dull the mower blade as well as create gaps in the sod. Moles also tunnel deep, throwing the excavated soil out of surface openings, thus forming molehills.

Practice Questions

1) Gradual metamorphosis has ____ developmental stages.

1. 1
2. 2
3. 3
4. 4

2) Complete metamorphosis has ____ developmental stages.

1. 1
2. 2
3. 3
4. 4

3) _____ feed by tearing plant cells and feed on sap.

1. Thrips
2. Caterpillars
3. Scale insects
4. Beetles

4) Wilting, puckering, or curling of plant foliage is a symptom of feeding by _____.

1. leaf miners
2. gall-makers
3. beetles
4. aphids

5) _____ are the stage of scale insects that are most susceptible to well-timed insecticide sprays.

1. Larvae
2. Crawlers
3. Pupae
4. Adults

6) Sooty mold is a fungus that grows in the honeydew produced by soft scales and some aphids.

1. True
2. False

7) The appearance of webbing and fine light spots on plant leaves during very dry conditions indicates an infestation of _____.

1. aphids
2. scales
3. spider mites
4. spiders

8) Only the adult stage of beetles are plant feeders.

1. True
2. False

9) Dogwood borers and lilac borers are caterpillars.

1. True
2. False

10) Skeletonizing is a type of leaf feeding that leaves a layer of leaf tissue and many veins undamaged.

1. True
2. False

11) The emerald ash borer only attacks very stressed or dying trees.

1. True
2. False

12) Most insect borers in Kentucky attack established, healthy trees and shrubs.

1. True
2. False

13) If large patches of dead turf are found late August that can be rolled up like loose carpet, the pest is probably _____.

1. sod webworms
2. armyworms
3. white grubs
4. cutworms

14) White grub control is the most effective way to control moles in turf.

1. True
2. False

Answers

- | | | | |
|-------|-------|-------|-------|
| 1: 3 | 2: 4 | 3: 1 | 4: 4 |
| 5: 2 | 6: 1 | 7: 3 | 8: 2 |
| 9: 1 | 10: 1 | 11: 2 | 12: 2 |
| 13: 3 | 14: 2 | | |

Diseases of Ornamentals and Turfgrass

Plant Disease Fundamentals

A **plant disease** is any harmful condition that affects a plant's appearance or function. Common pathogens that cause diseases include: fungi, bacteria, and viruses. Some nematodes are plant disease agents. Temperature extremes or nutrient deficiencies are examples of disorders caused by non-infectious factors.

Pathogens are fungi, fungus-like water molds, bacteria, viruses, and nematodes that cause infectious diseases of ornamentals and turf. They are microscopic. Plant disease pathogens may be spread in many ways:

- wind;
- rain;
- animals;
- soil;
- nursery grafts;
- vegetative propagation;
- contaminated equipment and tools;
- infected seed stock;
- pollen;
- dust storms;
- irrigation water; and
- people.

Infection begins when the pathogen enters the plant.

The disease process starts when it arrives at a part of a plant where infection can occur. If environmental conditions are favorable, the pathogen begins to develop. The plant is diseased when it responds.

Symptoms are the usually visible reactions of plants affected by these organisms. They may include leaf spots, chlorosis, cankers, galls, wilting, or root decay. Several types of pathogens can cause similar disease symptoms, so proper identification is very important.


Host-specific pathogens only infect certain hosts. If a host plant is not susceptible to infection, then the disease will not develop.

Also, **environmental conditions** must be right for infection. Most pathogens require wet conditions or high humidity so disease is usually much lower during very dry summers. On the other hand, rainy seasons usually produce more disease.

The **disease triangle** is a fundamental concept in plant pathology. **Disease occurs only when all three sides of the triangle are present: a susceptible host, a pathogen**

(the agent that causes disease), and an **environment favorable for disease to develop**. Plant diseases are managed by manipulating the disease triangle: the plant, the pathogen, and/or the environment.

The Plant Disease Triangle



Plants respond to disease in 3 main ways: **overdevelopment of tissue** - galls, swellings, or leaf curls; **underdevelopment of tissue** - stunting, lack of chlorophyll, or incomplete development of organs; or **tissue death** - blight, leaf spot, wilting, and cankers.

Sometimes, adverse growing conditions or environmental factors produce symptoms similar to those of plant diseases. These **abiotic problems (caused by non-living factors) need to be distinguished from plant diseases for proper management**. For example, fungicide applications cannot correct frost injury, dog urine burn, nutrient deficiencies, drought, girdling roots, changes in grade, chemical injury, air pollution injury, and mechanical damage.

Knowing the disease-causing organisms allows selection of the proper chemical or cultural practice to control the problem. For example, a root problem identified as a root rot when it is actually a nematode infestation will not be cured by applying a fungicide; a nematicide is required.

Accurate identification and diagnosis is an art, as well as a science, and experience is essential. This section will acquaint you with the general symptoms of diseases. For more accurate disease diagnosis, consult your county Extension agent.

Diagnosis of Plant Diseases




A correct diagnosis is the first step in disease management. You can recognize diseased plants by comparing them with healthy ones. To recognize a disease condition, you must know the plant's normal growth habits. When you are trying to identify the cause of a plant disease, you need to look for **symptoms** - the host plant's reaction to the disease agent, and **signs** - visible presence of the disease agent.

Many different plant diseases cause similar symptoms. Different pathogens and agents that are not pathogens can cause leaf spots, wilts, root galls, or stunted growth. For example, similar symptoms may be a result of mechanical injury, improperly applied fertilizers and pesticides, or frost. Often, the only way to pinpoint the cause is to find the observable signs that the particular disease agent is present -- such as fungal spores and mycelium or bacterial ooze.

Diseases of Ornamentals

Accurate identification and diagnosis of plant diseases is an art, as well as a science, and experience is essential. This section will acquaint you with the general symptoms of diseases of ornamentals. For more accurate disease diagnosis, consult your county Extension agent.



Fungi are multi-celled microbes that can either feed on living green plants or on dead organic matter. Disease occurs when pathogenic fungi attack living plants. Fungi usually produce spores which can cause infections when carried to a

susceptible plant. Spores can be moved by wind, water, insects, and tools.

Fungal spores require adequate moisture and the optimal air temperature in order to begin new infections. Many fungal diseases are common during wet, humid seasons. Some pathogens infect directly into healthy plant tissue, while others require a wound or other plant injury in order to invade plant tissue. Chemicals used to control fungi are fungicides. Usually, fungicides are applied to prevent, not cure, fungus disease.

Leaf Spots



Anthracnose of Maple (top) and Sycamore (bottom), photo: Univ. of Kentucky Nursery IPM



Black spot on rose, photo: Purdue University

Fungal leaf spots (such as anthracnose, scab, leaf blotch, or shot hole) can vary in size, shape, and color. Some spots have distinct margins and may be surrounded by

yellow halos. Other types of spots may be angular or blotchy. Spots or dead areas may enlarge to cover an entire leaf. As the spots become more abundant, leaves may yellow, die, and drop. Usually, leaf spots occur first on the lower leaves and progress up the plant. Fungal growth in the spot may consist of tiny pimple-like structures or a moldy growth of spores. You may need a hand lens or microscope to see the symptoms.

Leaf spots are more common in during early spring and fall when the moisture needed for infection is present. Fungal spores may overwinter in the infected leaves that drop around a plant. During spring, these pathogens produce infective spores that blow or splash onto healthy plants. If carried to healthy plants, these spores can begin a new infection under appropriate environmental conditions. Leaf spots occur on virtually all ornamental plants but not all leaf spot diseases affect plant health.

Leaf Blights



Dogwood anthracnose, photo: John Hartman, University of Kentucky, Bugwood.org

Leaf blights may have the same effect on plants as leaf spots but are **generally larger diseased areas and have less regular shapes.**

Dogwood anthracnose disease may begin as a leaf spot, become a leaf blight, and even progress to twigs and branches, causing dieback.

Rusts



Cedar-apple rust with telial "horns" (left) and cedar-quince rust (right), photo: Univ. of Georgia



Cedar apple rust on apple foliage and fruit photo: Univ. of Illinois

Rusts often produce spots called pustules that are similar to leaf spots. Pustules may be on the upper and/or lower leaf surface. They contain brown, reddish brown, orange, or yellow spores. Rust pustules are usually raised above the leaf surface. Rubbing the affected leaf surface will leave a dusty rust color (caused by the spores) on your fingers. Rust fungi may also attack twigs, branches, and fruit. They are often carried by wind and can be blown from infected plants to healthy plants, spreading the infection.

Rust diseases can have very complicated life cycles and, in many cases, require two separate hosts to complete their life cycle. In such cases, removing either one of the hosts can break the cycle and stop rust. Cedar-apple rust and related rusts are common ornamental disease problems.

Powdery Mildew



Dogwood powdery mildew, photo: UkNTrees

Powdery mildew The most common symptom is the white or gray layer of fungal growth produced on surfaces of the plant leaves and stems. Crooked stems or bubbled and curled leaves may develop if plant buds or very young tissue are infected.

Wind or rain splash can carry powdery mildew spores to new plants. During fall, the fungi produce small, black, overwintering structures that can overwinter in leaf debris or in cracks on bark.

Powdery mildew fungi are host specific, so different species infect different plant types. For example, powdery mildew fungi on dogwood will not infect Hydrangea, etc. Roses, oaks, tulip poplars, lilacs, zinnias, and euonymous are commonly affected by powdery mildew.

Leaf Gall Diseases



Leaf gall on lowbush blueberry photo: Bruce Watt, University of Maine, Bugwood.org

Leaf gall diseases are caused by fungi and are favored by cool, moist weather. However, most galls seen on plants are caused by insects or mites.

Leaf galls caused by fungi can usually be seen shortly after new growth begins in the spring. Parts of leaves become

distorted with a pale green to whitish bladder-like thickening. Young, thickened, fleshy leaves are covered with a white growth.

As galls age, they turn brown, dry up, and fall to the ground. If disease is severe, plant vigor can be affected due to leaf loss.

Dead, dry leaves that fall to the ground will be a source of spores for infection the following season. Leaf galls occur on azalea, camellia, and plum.

Root Rot



Phytophthora root rot, photo: John Ruter, University of Georgia, Bugwood.org

The first symptoms generally appear on above-ground plant parts as a gradual loss of vigor, yellowing of leaves, or wilting. Attempts to correct the problem with fertilizers and water generally yield little or no response.

In order to diagnose root rot diseases, plants must be dug carefully and soil washed from the roots. Diseased roots appear decayed, generally brown to black, and may be mushy or spongy.

The fungus-like water molds *Pythium* and *Phytophthora* and the fungi *Fusarium*, *Rhizoctonia*, and *Thielaviopsis* are common root rotting organisms.

Excess soil moisture favors root rot disease on ornamental plants. Once soilborne fungi build up in landscapes, it is difficult to disinfest soil.

Stem Rot / Stem Blight



Stem rot, photo: Matt Montgomery, Sangamon-Menard Extension

The pathogens commonly associated with stem rot of ornamentals include the fungus-like water molds *Pythium* and *Phytophthora*, and the fungi *Rhizoctonia*, *Sclerotium*, and *Botrytis*. All are common soil-inhabiting fungi.

They can be spread in infected debris, on cuttings, or when soil is moved. Once soilborne fungi build up in landscapes, it is difficult to disinfect soil.

Plants infected with stem rot fungi often show early symptoms of wilt. During advanced stages of disease, plants become more severely wilted and eventually die. The stems may be brown and shrunk at the soil line. Under extremely moist conditions, the white, cottony fungus mycelium may be visible on the surface of the stem.

Chrysanthemums, geraniums, petunias, and other herbaceous ornamental plants are very susceptible to stem rot.

Damping-off, a similar disease of seedlings, kills ornamental seedlings during the first few weeks after seed germination.

Cankers



Stem canker on Japanese rose, photo: Elizabeth Bush, Virginia Polytechnic Institute and State University, Bugwood.org



Canker on Walnut, photo: Paul A. Mistretta, USDA Forest Service, Bugwood.org

Cankers are localized sunken lesions areas on trunks, stems, or branches of woody plants. Canker diseases cause bark tissues to shrink and die. The dead tissues often crack open and expose the wood underneath.

Cankers begin as small, discolored yellow, brown, or red spots that sometimes appear water-soaked, although some canker disease are not visible outside of the bark. As cankers enlarge, their centers may become tan or gray. Small, black, pimple-like structures (fungal fruiting bodies that contain spores) may form in the canker. Cankers can enlarge and girdle stems, causing death to parts of the plant above the canker.

Fungi causing cankers usually infect through a wound or injury to the bark or wood. Rose canker is a common example of a disease showing this symptom.

Vascular Wilt Diseases



Verticillium dahliae fungus causes this vascular wilt disease on hundreds of woody plants photo: JW Pscheidt, U Mass.

Fungal pathogens such as *Fusarium*, *Verticillium*, and *Ophiostoma* can cause wilting of many ornamental species by restricting the water flow to leaves and stems. The wilting caused by such pathogens is sometimes due to the toxins they produce. Other pathogens can build up within water-conducting vessels, which become plugged by fungal growth.

Vascular wilt diseases often affect one side of the plant first, causing individual limbs or branches to wilt and die back. *Fusarium* and *Verticillium* infections usually begin in roots and gradually spread internally throughout the infected plant. *Verticillium* wilt of maple is an example.

Other wilt fungi infect through upper plant parts. Symptoms of vascular wilt disease often include discolored streaks in the wood of infected branches, which are visible upon cross sectioning infected wood.

Bacteria



Bacterial leaf scorch, photo: John Hartman, University of Kentucky, Bugwood.org

Bacteria are single-celled organisms that usually reproduce by simple cell division, some as often as every 30 minutes. They can build up quickly under warm, humid weather conditions. Leaf, growing shoots, and fruit diseases are the most common types in Kentucky.

Bacteria can be carried from plant to plant in water droplets, by wind, rain splash, insects, or on equipment. They often survive between growing seasons in crop residue, in seeds or cuttings, or in weeds.

Viruses



Rose mosaic virus, photo: William M. Brown Jr., Bugwood.org

Viruses are too small to see with a microscope. Generally, they are recognized by their effects on plants. These

include stunted growth; change in plant color; abnormal formation of infected roots, stems, leaves, or fruit. Mosaic diseases, characterized by light and dark blotchy patterns, usually are caused by viruses.

It can be difficult to distinguish between diseases caused by viruses and those caused by other plant disease agents, such as fungi and bacteria. A positive diagnosis requires sophisticated testing, such as inoculating indicator plants and observing the results or using specifically identified antibodies to test for the presence of the organism.

Viruses depend upon living organisms for food and reproduction; they cannot exist very long outside a host. They are commonly spread from plant to plant by mites, aphids, leafhoppers, or whiteflies. A few are spread in the seeds of the infected plant.

Nematodes



Foliar nematode damage on Hosta photo: Department of Plant Pathology, North Carolina State University, Bugwood.org

The life cycle of a nematode includes an egg, several larval stages, and an adult. Most larvae look like small adults.

In adverse conditions, females of some species form inactive, resistant forms called **cysts**. The cyst is the hard, leathery, egg-filled body of the dead female, which is difficult to penetrate with pesticides. A cyst may protect eggs for as long as 10 years.

Diseases of Turfgrass

Accurate identification and diagnosis of plant diseases is an art, as well as a science, and experience is essential. This section will acquaint you with the general symptoms of diseases of turfgrass. For more accurate disease diagnosis, consult your county Extension agent.

Numerous disease problems occur on turfgrass in Kentucky and these frequently cause extensive damage.

In many cases, a disease is blamed for poor quality turf when, in reality, it may be only a contributing factor or not involved at all. Frequently, dead and dying grass is caused by improper fertilization, chemical burn, mower problems, dog or insect injury, dry or wet spots, thatch, competition from other plants, or from any other improper management. Accurate diagnosis of the problem is essential for proper control.

Two types of pathogens (fungi and nematodes) are found in turf in Kentucky. Observation of symptoms is an important aid in determining which pathogen is causing a disease. The following information will explain the identification and biology of some common turf diseases.

Helminthosporium Leaf Spot



"Football" shaped lesions of leaf spot and melting out of Kentucky bluegrass photo: S. Tirpak, Rutgers

Helminthosporium leaf spot is a common disease problem of Kentucky bluegrass and is often referred to as "melting-out." Other *Helminthosporium* leaf spots are important on fescues and bermudagrass. From a distance, leaf spot-affected areas appear chlorotic or yellowed. Individual spots on the leaves have dark margins with tan centers. The spotting is most noticeable in spring and early summer.

Infection in the crown of the plant during the summer can lead to the death of plants (thus "melting-out"). Cool, wet weather during spring followed by drought during summer accentuates the damage from this disease.

Dollar Spot



Dollar spot, photo: Michigan State Plant Pathology

Dollar spot appears as sunken round, bleached-out or straw-colored spots and affects a wide variety of grasses, including Kentucky bluegrass, bermuda grass, perennial ryegrass, zoysia, tall fescue, and bentgrasses. The fungus is active throughout the growing season, especially when there is low soil moisture and an excess of dew or fog. It is most prevalent in the spring.

The disease is characterized by small white patches, one to three inches in diameter. A large number of spots can come together and form larger dead areas. Leaf spots are usually found along the edges of the grass blade and may come together across the blade, causing the tip to die. Individual leaf spots are tan with reddish margins.

Pythium Blight



Pythium blight, photo: William M. Brown, Jr., Bugwood.org

Pythium blight is caused by a number of species of the fungus *Pythium*. The fungus primarily attacks perennial ryegrass and bentgrass although other grasses can be affected. Conditions that favor *Pythium* blight include abundant moisture and poor air circulation. The disease is most active in hot, humid weather when the night temperature does not go below 70°F.

The blight appears first as small spots a few inches in diameter. Diseased leaves are at first water-soaked, soft, and slimy, and may mat together. Dense, cottony fungal growths often are apparent in affected areas during a heavy dew. The leaves soon shrivel and the color of the patch soon fades to light brown as dew dries. The shape of the diseased area may be streaked following the drainage flow of water over the turf.

Brown Patch



Brown patch on bentgrass, photo: University of Massachusetts

Brown patch is a common fungal disease of fescues, perennial ryegrass and bentgrass. It develops most readily when daytime highs exceed 80°F and nighttime lows are in the mid- 60's°F or higher.

Brown patch is one of the more common turf diseases, especially in tall fescue. In addition to ideal temperatures and humid weather, heavy applications of nitrogen fertilizer favor disease development.

Brown patch is characterized by nearly circular areas of dead leaves that may be a few inches to several feet in diameter. On closely mown turf, the edges of the dead area may have a gray, smoky color, particularly in early morning. Affected areas are generally tan or brownish in bent and ryegrass. Affected fescues usually have straw-colored leaves.

Summer Patch



Summer patch, photo: Mary Ann Hansen, Virginia Polytechnic Institute and State University, Bugwood.org

Summer patch affects Kentucky bluegrass and annual bluegrass. Circular to irregular patches of dead turf up to 1-2 feet in diameter develop during hot weather in mid-to late summer. Below ground, roots and crowns of affected plants are brown and decayed, a result of fungal colonization. A tuft of healthy, green grass is sometimes evident in the center of affected patches, giving them a characteristic “donut” appearance.

Necrotic Ring Spot



Necrotic ring spot, photo: Howard F. Schwartz, Colorado State University, Bugwood.org

Necrotic ring spot is another disease of bluegrasses with symptoms similar to summer patch. In contrast to summer patch, symptoms of necrotic ring spot can develop following cool, wet weather in late spring or mid-autumn. Necrotic ring spot is less common in Kentucky than summer patch.

Rust



Rust pustules on turfgrass, photo: Iowa State University

Rust is sometimes a problem on Kentucky bluegrass, fescue, zoysia, perennial ryegrass, and Bermuda grasses. Rust infection results from rust spores which are blown to the plant from distant areas or from nearby alternate hosts. Large numbers of spores are produced in the leaf spot (pustule). These spores are then the source of new infections.

The disease is most frequently found during cool, humid weather during autumn. Grass varieties differ in susceptibility to rust.

Red Thread



Red thread, Bruce Watt, University of Maine, Bugwood.org

Red thread is seen as irregularly shaped patches of blighted turfgrass, ranging from a few inches to a few feet in diameter. Often, as diseased leaves turn brown, pink or reddish fungal growth can be observed on the leaf surface or emerging from the cut ends of leaves. This disease affects most of the common grasses grown in Kentucky and is often found during spring and early summer. The disease is favored by conditions of low nitrogen fertility.

Mushroom Fairy Ring



Mushroom fairy ring, photo: Clemson University - USDA Cooperative Extension Slide Series, Bugwood.org

Mushroom fairy ring can occur in any turf. The ring appears as a circular discoloration of grass from several inches to many yards in diameter. Mushrooms (toadstools) may appear at the edge of the ring during warm, moist periods).

The ring of grass is generally a darker green than the grass inside and outside the ring. During periods of moisture stress, the grass inside the ring may die. This general decline of grass inside the ring adds to the unsightliness of the fairy ring problem. Fairy rings gradually increase in size.

Slime Molds



Dog vomit slime mold, photo: Sandra Jensen, Cornell University, Bugwood.org

Slime molds are commonly found on lawns in warm, moist weather. This fungal growth on grass leaves may be either a small, crust-like, light to dark mass with a sooty appearance, or a tan to orange shapeless mass.

The fungus causing this unsightly problem does not infect the grass blade; it simply uses it for support. The only

effect it has on the plant is to temporarily reduce food production by the grass leaf as a result of shading.

Nematodes



photo: Bonsak Hammeraas, NIBIO - The Norwegian Institute of Bioeconomy Research, Bugwood.org

Root-gall nematodes on Kentucky bluegrass

Nematodes weaken and reduce the vigor of turfgrass by restricting the development of the root system. The symptoms of nematode injury may be confused with nutritional problems, insufficient water, hardpan, or any factor that restricts root development.

Symptoms commonly associated with nematode injury include thinned or completely killed areas, pale green to chlorotic color, excessive wilting during drought stress, poor response to fertilization, and a greater weed problem due to sparse grass.

The intensity of the symptoms will vary with the grass variety, the kinds of nematodes present, the nematode population level, and the fertilization-watering program being practiced. The most reliable method for determining whether a nematode problem exists is by a soil assay. Nematode damage to turfgrass is uncommon in Kentucky.

Practice Questions

1) A _____ is a rapid discoloration and death of tree foliage.

1. canker
2. gall
3. blight
4. dieback

2) _____ is an abnormal loss of normal green leaf color.

1. Distortion
2. Gall
3. Defoliation
4. Chlorosis

3) _____ is a burning that moves into the leaf from its margin.

1. Gummosis
2. Leaf scorch
3. Leaf spot
4. Wilt

4) _____ are very dependent upon water to spread from plant to plant.

1. Bacteria
2. Fungi
3. Viruses

5) _____ are identified by mycelia, spores, and fruiting structures.

1. Bacteria
2. Nematodes
3. Fungi
4. Viruses

6) Many plants respond to oxygen starvation by dropping the lower leaves that are usually yellowed or necrotic.

1. True
2. False

7) Which of the following causes a plant disorder?

1. fungus
2. bacteria
3. nematode
4. nutrient deficiency

8) _____ are the most common infectious organisms causing plant disease.

1. Bacteria
2. Fungi
3. Sunscalds
4. Viruses

9) _____ reproduce by spores.

1. Nematodes
2. Bacteria
3. Viruses
4. Fungi

10) _____ reproduce by simple cell division.

1. Bacteria
2. Fungi
3. Nematodes
4. Viruses

11) _____ is an example of a fungal disease.

1. Root cyst
2. Mosaic patterns on leaves
3. Powdery mildew
4. Leaf blight

12) Wilting is an example of a disease sign.

1. True
2. False

13) An infection begins when the pathogen _____.

1. enters the plant
2. arrives at a part of the plant where infection can occur
3. symptoms appear

14) _____ is the first step in disease management.

1. Waiting for symptoms to appear
2. A correct diagnosis
3. Selection of the most effective fungicide

Answers

1: 3 2: 4 3: 2

4: 1 5: 3 6: 1

7: 4 8: 2 9: 4

10: 1 11: 3 12: 2

13: 1 14: 2

Pesticide Application Equipment and Methods

The application method you choose depends on such factors as the nature and habits of the target pest, characteristics of the target site, and properties of the pesticide formulation. You also must consider the suitability of the application equipment, cost, and efficiency of alternative methods.

Here are some common pesticide application methods:

Band Application



These are often applied along fence rows or borders with a non-selective herbicide to kill all vegetation.

Broadcast Application



Pesticides are spread uniformly over a large area of turfgrass on foot or with motorized equipment.

Drench Application



These are used to treat specific areas usually with systemic products that are taken up by the roots and moved throughout the plant.

Foliar Sprays



These sprays are directed toward the leafy portions of the plant.

Soil Applications



Pesticides are placed directly on or in the soil instead of on the plant.

Space Treatment



Pesticides are applied to the air inside of an enclosed space.

Spot Treatment



This is an effective way to treat specific problem areas without treating the entire turf area. Be careful not to walk through the treated areas when using non-selective

herbicides like glyphosate and watch for dripping from the nozzle while walking from site to site.

Wiper Application



This can be used with a non-selective herbicide to selectively kill individual weeds. The wiper's wetness must be less than would cause dripping and handled carefully to avoid accidentally touching desired plants.

Safety Systems

Pesticide Containment Pad

If you often store, handle, mix and load pesticides, or clean equipment at the same location, you may have to install a pesticide containment pad. Check EPA and Kentucky state regulations to determine when a containment pad is required. These pads are designed to contain spills, leaks, overflows, and wastewater for reuse by the applicator or for disposal by a commercial waste management contractor. They make it easier to clean up spills and help to prevent environmental contamination.



Impervious containment pad

Generally, the containment pad **must be made of impermeable material**. It should be **concave or have curbs, berms, or walls high enough to hold the largest amount of spill, leak, or equipment wash water** likely to occur at the site. It also must have a **system to remove and recover spilled, leaked, or released material** by

either an automatic sump system or a manually operated pump. Smaller, portable pads and lightweight trays made of heavy-duty plastic may be used when mixing and loading at the application site.

Hydraulic Sprayers

The application equipment or device **must be able to apply the pesticide to the intended target at the proper rate**. The label specifies the legal application rate and may suggest the appropriate equipment for use with the product.

Hydraulic sprayers range from powered units with a multiple-nozzle boom to a handpumped backpack sprayer. In all cases, pressure from either a pump or compressed gas or air is used to atomize the spray mix at the nozzle. High pressure pumps are needed to provide good spray coverage on large trees. Manual sprayers are designed for spot treatments and for areas unsuitable for larger units. They are relatively inexpensive, simple to operate and maneuver, and easy to clean and store. Adjustable spray guns are often used for lawn care sprays.



Broadcast sprayer (BASF)



Backpack sprayer (Solo)



Lawn spray (Mid Dept Ag)



Tree sprayer (Masspray.com)

Sprayer Components



Tank

A **tank** is necessary to contain the spray mix. **Choose one made of, or coated with, a material that does not corrode and that can be cleaned easily.** Cleaning prevents accumulations of corrosion and dirt that clog screens and nozzles, increasing wear on the equipment. Large tanks require an opening in the bottom to aid in cleaning and draining. A large top opening is useful for filling, cleaning, and inspecting the tank. The opening must have a watertight cover to prevent spills. A **tank agitation system/device** is useful for most sprayable formulations, especially for wettable powders or dry flowables. Constant mixing of a pesticide and liquid carrier produces a uniform spray mixture (suspension) and results in an even application of the chemical.

Exposure to sunlight and corrosive chemicals can shorten the life of polyethylene tanks.

Three common signs of wear and potential tank failure are:




photo: omafra.gov.ca

- **Scratches** are on the surface and can be seen and felt
- **Crazing** is a network of fine lines or cracks that may look like a patchwork, but often cannot be seen with a visual inspection. Crazing can be seen when using one of the testing methods explained below. **Crazing occurs within the tank wall and can be a sign of deterioration of the plastic, which may lead to cracks.** Tanks that show signs of crazing will still hold liquids, but the integrity of the tank is questionable. For this reason, caution should be used when putting any hazardous substance in tanks that show crazing.
- **Cracks** extend through the plastic wall and can be visually seen and felt. Cracks may run parallel or at right angles to each other.

Pump

A **pump** agitates the spray mixture and produces a steady flow to the nozzles. Pump parts must resist corrosion and abrasion, especially when wettable powders or similar formulations are used. Never operate a sprayer pump at speeds or pressures above those recommended by the manufacturer. You may damage the pump if it is operated dry or with a restricted flow at the inlet or outlet. Pumps depend on the spray liquid for lubrication and to prevent overheating.

Nozzles

The proper selection of a **nozzle** type and size is essential for proper pesticide application. **The nozzle is a major factor in determining the amount of spray applied to an area, the uniformity of application, the coverage obtained on the target surface, and the amount of potential drift.**

Nozzles break the liquid into droplets, form the spray pattern, and propel the droplets in the proper direction. Nozzles determine the amount of spray volume at a given operating pressure, travel speed, and spacing. Drift can be minimized by selecting nozzles that produce the largest droplet size while providing adequate coverage at the intended application rate and pressure.





Photo: TeeJet.com

- 1) The **nozzle body** holds the strainer and tip;
- 2) The **strainer** screen prevents a clogged nozzle. It is the best defense against nozzle plugging and pump wear. The screen can remove dirt and rust flakes from the spray liquid before it reaches the nozzle.
- 3) **Tip gasket** to reduce leaks and dripping.
- 4) The **spray tip** determines the flow rate and droplet pattern;
- 5) The **cap** holds the nozzle body and tip in place.

The Spray Tip

The **spray tip** determines the flow rate and droplet pattern.



11004 nozzle


110 is the spray angle in degrees, 04 is the output - 0.4 gallons per minute at 40 psi, photo source: TeeJet




Equip nozzle tips with **check valves** to help prevent dripping when the pump is off. Be sure the spring-loaded ball valves are working properly.

Nozzle Spray Angle, Spacing, and Boom Height

Nozzle spray angle is formed by the edges of the spray pattern. Common angles are 65°, 80°, and 120°. A wide-angle nozzle (110°) produces a thinner sheet of water with smaller droplets than a narrow angle nozzle (65°) with the same delivery rate.




Nozzle spacing on the boom, spray angle, and boom height determine proper overlap of the spray. The drawing below shows the effect of nozzle spray angle on nozzle height, need for proper overlap, and spray coverage. Notice the height difference between the 110°, 80°, and 65° nozzles. Wide angle nozzles are placed closer to the target for proper overlap. A lower nozzle height reduces the risk of spray drift.



Flow meters and other devices measure the uniformity of nozzle flow rate from nozzles along a boom. They are very useful when calibrating sprayers with multiple nozzles.

Common Nozzle Spray Patterns



Three common nozzle spray patterns: hollow cone, full cone and flat fan, photo: sprayers101.com

Hollow cone nozzles produce a fine spray pattern to completely cover leaf surfaces. **Full cone nozzles** produce large, evenly distributed droplets at high flow rates. These two cone nozzles are best suited to apply fungicides and insecticides. **Flat fan nozzles** form narrow, oval patterns

with tapered ends. They are spaced along a boom and overlap by 30% to 50% for even broadcast spray distribution to the soil surface or plant canopy.

Nozzle Maintenance



Nozzles are available in various materials: brass, aluminum, plastic, stainless steel, hardened stainless steel, and ceramic. Select the material best suited for the pesticide formulation being used.

Never use brass or aluminum tips to apply abrasive materials (such as wettable powders and dry flowables) because they wear too fast. This wear increases the opening size of the nozzle, which increases its output. Reduce wear by using nozzle tips made of a hard, wear-resistant material: plastic, hardened stainless steel, or ceramics.

Be sure you have the correct screen size for each nozzle.

Clean nozzle tips carefully with a soft brush, not wire or a knife tip.

Sprayer Cleanup

Spray equipment should be cleaned in the field after the spray job has been completed. Some pesticide labels provide specific information on cleaning spray equipment; consult the label for guidelines. **Do not clean spray equipment in areas where rinse water will contaminate water supplies, streams, or injure susceptible plants.**

Pay special attention to areas that can be missed or are difficult to clean:

- Spray surfaces or components **where buildup of dried pesticides might occur**
- Sprayer **sumps and pumps**
- **Inside the top of the spray tank** and **around baffles**
- **Irregular surfaces** inside tanks caused by baffles
- **Plumbing fixtures**, agitation units, etc.
- **Collection points** where the hoses connect to the nozzle fittings in dry boom sprayers.

Flushing spray equipment with water may be sufficient to remove potentially harmful amounts of many pesticides. However, certain groups of pesticides may require special attention. **Thorough clean-out procedures can be critically important when switching applications between crops to help avoid significant crop injury.** As a rule, a sprayer that has been used to apply 2,4-D or other growth regulator type herbicides should not be used to treat susceptible plants. A triple rinse – water, then ammonia, then water again – minimizes the risk of injury from dicamba and 2,4-D.

Types of Sprayers



The **backpack sprayer** is a simple but useful piece of application equipment made up of a tank, pump, spray wand and nozzles. It is useful for treating small areas, spot sprays, or hard to reach locations. The main spray options are broadcast, band, and spot.

Broadcast and band spraying are used to treat areas uniformly. They require determining spray pressure, walking speed, nozzle tip, and height. Spray pressure is maintained by hand pumping. Determine a comfortable constant walking speed for the slope and terrain you will be covering. Select a nozzle tip for the volume of mixed spray that you will be applying. In many cases, the spray rate is given in teaspoons or ounces per 1,000 square feet.

Spot spraying is common for treating scattered clumps of weeds or brush. In many cases, you mix a specific concentration, such as a tablespoon per gallon of water and apply it until the foliage is wet but not dripping.

Hose reel lawn care sprayers allow efficient, even application of pesticides and fertilizers to turf. However, practice, experience, and attention to walking speed and application technique are essential for effective applications.



photo source: Turfsprayers.com

Common outputs for spray gun nozzles are 1.5, 3, and 4 gallons per minute.



photo source: krittenhouse.com

Inconsistent applications can result in control failures or injury to turf or landscape plants. Things to watch for when using this type of application:

- Pump pressure is set correctly
- Consistent, accurate walking speed
- Nozzle is correct for desired flow rate
- No partially blocked nozzle openings
- Hose is not kinked
- Strainer screen is not clogged

Granular Applicators




Rotary and drop spreaders, Penn State Univ.

Rotary and drop spreaders are used to apply granular fertilizers and pesticides to turf. The spreaders have several holes in the bottom of the hoppers with moveable gates which can be set to regulate product flow. Gate

openings are adjusted during the calibration process to give the proper delivery rate.

The swath of the rotary spreader (above) is 12 feet but the distribution of the granules from it is uneven. You must overlap swaths to get an even application. The amount of overlap varies among spreaders. You need to measure the “effective swath width” of your spreader to determine the amount of overlap that is needed for uniform coverage.



Uneven distribution of granules from a rotary spreader, photo: Pesticide Environmental stewardship - CIPM

Lawn care products applied by drop spreaders fall directly beneath the equipment. Leaving a gap between swaths can result in streaking while overlap will double the rate.



Uneven fertilizer application: gaps between passes with a drop spreader, photo: Donnan.com

Non-spray Alternatives to Treating Trees

(source: modified from Utah State Univ. pub NR/FF/020)

Pesticides can be applied to tree trunks, branches, and foliage. This approach has been used extensively in recent years to protect ash trees from the emerald ash borer. Differences in chemical characteristics of products and advances in alternative application methods allow trunk implantation, trunk injection, soil injection/drenching, and trunk basal sprays. **These approaches allow more efficient use of pesticides, more effective placement, especially against some borers, and elimination of drift. However, movement of some chemicals in to nectar and/or pollen can cause significant harm to pollinators.**

Trunk Implants



photo source: Gardenersedge.com

Trunk implants or injections work by placing water soluble pesticides at or in the cambium where they can be carried through the tree mainly in xylem sap. The **pesticide must be placed in the correct tree tissue at the right time of year and at an effective concentration.** This tends to be a **good way to treat many sap feeding insects, borers, and some caterpillars.** Implantation involves placing capsules of pesticide into the outer xylem or sapwood. The product coating is dissolved by transpiration water and the chemical moved to the target site. Implants tend to require relatively large holes.

Trunk Injection

This involves placing pesticides into the tree for direct uptake. **Injections are necessary for large trees and can be used on sites where soil treatments may not be practical, effective or appropriate,** including trees growing on excessively wet, sandy, compacted or restricted soil environments. Trunk injections generally

involve drilling through the bark and into the outer sapwood at the base of the tree. Drilling wounds could cause long-term damage, especially if treatments are applied annually.



Setting up a trunk injection, photo: moorparktreeservice.com

Application methods that rely on high pressure injections of insecticide through needles inserted into small holes **may damage the tree if the pressure causes the bark to bulge and separate from the cambium.** This is most likely to occur in spring and can cause larger wounds that result from death of the vascular tissue at the point of separation.

Uptake of trunk-injected insecticides will be most efficient when trees are actively transpiring. Best results are usually obtained by injecting trees in the morning when soil is moist but not saturated. Uptake will be slowed by hot afternoon temperatures and dry soil conditions. Irrigating trees during droughty conditions will help with insecticide uptake and translocation within the tree. Products applied as trunk injections are typically absorbed and transported within the tree more quickly than soil applications. **Allow at least two and preferably three to four weeks for most trunk-injected products to move through the tree.**

Soil Injection

Soil injection relies on placing the pesticide, diluted in water, into the soil for uptake by the root system of the tree. While no holes need to be drilled, as with implants or injections, these applications generally **must be made several weeks before the pest is active to allow time for uptake and dispersal of the chemical into the tree.** In general, there is a limit to the size (diameter) of trees that can be treated by this procedure. They require specialized equipment, but offer the advantage of placing the insecticide below mulch or turf and directly into the root zone of the tree. This also can help to prevent runoff on slopes. **Injections should be made just deep enough to place the insecticide beneath the soil surface (2-4 inches).** Soil injections should be made within 18 inches of the trunk. Studies have shown uptake is higher and the

treatment more effective when the product is applied at the base of the trunk where the density of fine roots is highest.



Hand operated soil injection device, photo: Virginia Tech University

No soil applications of systemic insecticide should be made where there are roots of flowering plants that are visited by bees and other pollinators. This situation is most likely to occur where flowering plants are established around the base of an ash tree. In these situations the flowering plants should either be destroyed or insecticide should be applied via trunk injection to ensure the toxins will not be taken up by the flowering plants.

Trunk Basal Spray

The **trunk basal spray** alternative is based on thoroughly wetting the lower 5 feet of the trunk with a water soluble pesticide. The chemical is absorbed through the bark and distributed by the vascular system of the tree. The basal trunk spray offers the advantage of being **quick and easy to apply and requires no special equipment** other than a garden sprayer. This application technique does not wound the tree, and when applied correctly, the insecticide does not enter the soil. Sprayers must be calibrated to ensure the appropriate amount of the formulated product is applied to each tree.



Photo source: utahpests.usu.edu

There are advantages and disadvantages to each alternative but **all require largely intact vascular systems to move the pesticide.**

Practice Questions

1) The nozzle _____ determines the flow rate and droplet pattern.

1. body
2. tip
3. cap
4. strainer

2) Drift can be minimized by selecting a nozzle that produce the smallest possible droplet size.


1. True
2. False

3) The strainer screen in a nozzle assembly _____.

1. increases spray pressure
2. stops dripping during turnaround
3. eliminates spray drift
4. keeps the nozzle tip from clogging

4) The spray angle and flow rate of this nozzle is _____ degrees and _____ gallons per minute.

1. 800 and 4
2. 40 and 0.8
3. 80 and 0.4
4. 80 and 4



5) Hollow cone and full cone spray nozzles are best suited to apply pesticides evenly to the soil surface.

1. True
2. False

6) Strainer screens ARE NOT needed to catch product particles when applying only _____ formulations.

1. DF
2. WDG
3. F
4. EC

7) Which nozzle material is the best choice when applying mostly WP, DF, or WDG formulations?

1. aluminum
2. stainless steel
3. brass

8) Clean a clogged nozzle tip with a _____.

1. soft brush
2. knife blade tip
3. piece of stiff wire

9) A check valve will help to _____.

1. keep the nozzle flow steady if the pressure drops
2. reduce drift
3. calibrate the nozzle
4. prevent dripping when the pump is off

10) Flushing spray equipment with water is an effective way to remove residues of growth regulator herbicides.

1. True
2. False

11) Switching between granular products with smaller OR larger or lighter or heavier granules requires adjusting the flow rate of a rotary spreader.

1. True
2. False

12) A(n) _____ application of a herbicide is the method used to kill weeds along a fence row.

1. broadcast
2. drench
3. injection
4. banded

13) Common outputs for hose reel lawn care sprayers are _____ gallons per minute.

1. 1.5 to 4
2. 10 to 15
3. 20 to 25



14) A network of fine lines or cracks on polyethylene spray tanks indicates _____ (See above).

1. a strong tank agitator
2. residues of spreader stickers
3. cracking due to excessive sprayer pressure
4. crazing due to sunlight and corrosive chemicals

15) Loose gravel is the best type of material to use for a pesticide containment pad.

1. True
2. False

16) _____ spraying is a common way to treat scattered clumps of weeds or localized groups of pest insects.

1. Broadcast
2. Spot
3. Banded
4. Random

17) Which spreader will give a more even application of a mix of different sized or weight granules?

1. Hydraulic
2. rotary
3. drop

18) Three 30- to 60-second rinses of a sprayer with cold water minimizes the risk of injury from residues of growth regulator herbicides (2,4-D, etc.) in the tank and hoses.

1. True
2. False

19) The distribution of granules from a rotary spreader is very even across the entire swath width so no overlap is needed.

1. True
2. False

20) Trunk injections of pesticides may be necessary for _____ trees.

1. deciduous
2. softwood
3. large diameter
4. small diameter

21) Uptake and movement of trunk-injected pesticides will be most efficient when trees are dormant.

1. True
2. False

Answers

1: 2	2: 2	3: 4	4: 3
5: 2	6: 4	7: 2	8: 1
9: 4	10: 2	11: 1	12: 4
13: 1	14: 4	15: 2	16: 2
17: 3	18: 1	19: 2	20: 4
21: 2			

Potential Consequences of Pesticide Use

Pesticides can be very important and effective tools to manage specific problems, but applying pesticides in areas with diverse landscapes and around homes where children play and pets roam may increase the risk of unintended consequences. While these concerns are not unique to turf, lawn, ornamental or interior landscape applications, potential problems due to drift, phytotoxicity, persistence, and effect on non-target organisms must be managed.

Drift

Pesticide spray drift is the off-target movement of pesticide dust, droplets, or vapors through the air at the time of application or soon after, out of the intended area. Pesticide droplets are produced by spray nozzles used in application equipment. Pesticide drift can affect people's health, the environment, and damage nearby plants. Drift can result in less effective control as less material remains on target. Some products and formulations are more prone to drift, but to a limited degree there is a degree of drift associated with nearly all pesticide applications. For these reasons, drift should be considered and managed with all pesticide applications. Many factors can affect drift and damage caused by drift including weather conditions (wind speed and direction, humidity, temperature), type of pesticide and its formulation, distance to sensitive areas, spraying techniques (nozzle type, pressure, carrier volume, nozzle height, etc.), and applicator decisions. Drift can lead to liability issues for your business and support arguments to ban some pesticides.

Situational Awareness

Before making an application, understand what and where are the sensitive areas near the area to be treated. These can include, commercial crops as well as vegetable gardens, fish ponds, pollinator gardens and other flowering plants, bee hives, streams, and storm drains. Avoid making applications when prevailing winds will favor drift toward these areas.

Applications of granular pesticide that land on impervious surfaces like sidewalks and driveways can result in runoff into bodies of water. In these instances, sweeping these products off of these surfaces will reduce the chance of off-site movement.

Weather Conditions

Wind speed is a critical factor that moves pesticide dusts and droplets out of the intended areas. Some labels have specific restrictions on use based on wind speed, so applicators must be aware of those restrictions on the labeling. For those without specific restrictions, wind speeds of between 3 and 10 mph in a safe direction are recommended. Wind direction away from nearby sensitive areas can greatly reduce the risk of harm to sensitive areas, but wind direction should be used as a reason to apply pesticides when conditions favor excessive drift.

Drift and Pesticide Product Type

Some products are more prone to drift, and some products are more prone to cause damage when they do drift. Some of the auxin mimicking herbicides are prone to volatilize and move off-site. Non-selective herbicides are more likely to damage surrounding vegetation when they drift. The risk is elevated in plantings with a diversity of plant types and species.

Drift Reduction Technology

One rule of drift is that small droplets travel much farther than large droplets. To increase droplet size, applicators can increase the size of nozzles to deliver more spray volume per area, reduce sprayer pressure and speed of applications, or switch to drift reduction nozzles. Each of these techniques will require that the equipment be recalibrated for each change in nozzle type and size, and spray pressure.

Phytotoxicity and Persistence

Both phytotoxicity and persistence can be more of a concern in areas with a high diversity of plants. Persistence of some pesticides can result in phytotoxicity to some sensitive plants due to pesticide residues remaining in the soil as plantings are changed over time. Phytotoxicity can also be the result of drift. Phytotoxicity can take many forms and may cause a delay of seed germination, inhibition of plant growth or any adverse effect on plants including deformed growth, chlorosis, spotting, marginal burn, or tissue death.

In interior plantscapes, plants are often located adjoining large glass areas, plant damage can be minimized when pesticides are applied during the cooler part of the day. Avoid applications during hot, sunny weather or when the plants are stressed.

Spray tank mixtures of insecticides, miticides or fungicides may result in plant injury that does not occur from use of any one of the materials alone. Before materials are tank mixed, study the manufacturer's label carefully. Mixing pesticides that require different types of adjuvants should be avoided. It is best to treat just a few plants with a new combination of pesticides and wait a week for any phytotoxic effects to appear.

Non-Target Effects to Other Organisms

Turfgrass and lawns are recreational areas for people and companion animals and foraging areas for wildlife. Always read and following all label directions and restrictions. Application equipment should be calibrated regularly to ensure the dosages applied are within the labeled range. Children's toys and food preparation items should be removed or covered as needed. Keep children, adults, and pets out of treated areas as required by the label or until sprays dry, and posting and notifying customers and neighboring residents as required to ensure applications are conducted properly.

Pollinator Protection

Kentucky has a pollinator protection plan that is available [online](#) that addresses best management practices for pesticide applicators. Insect pollinators are particularly sensitive to some insecticides when they drift to their nests or contaminate flowers that they pollinate. Many native trees, shrubs, and wildflowers are dependent on insect pollinators in addition to the agricultural crops that rely on them.

Pesticide Applicator Best Management Practices:

- Be aware of honey bee hives and habitat for other pollinators near areas to be treated.
- Use IPM and thresholds when available for making application decisions.

- Avoid using dusts and wettable powders in pollinator-sensitive areas.
- Consider impacts on pollinators when making pesticide application decisions.
- Always use pesticides according to the label and follow all pollinator protection restrictions.
- Minimize pesticide drift and avoid applications when weather conditions favor pesticide movement toward honey bee hives.
- Notify nearby beekeepers prior to application if required by the pesticide labelling.

Practice Questions

1) Granular pesticides that land on impervious surfaces like sidewalks and driveways are not a problem as they can be washed into the storm sewer.

1. True
2. False

2) Small droplets travel much farther than small droplets with the wind.

1. True
2. False

3) Which of the following can affect pesticide drift?

1. Weather conditions
2. Type of pesticide
3. Application techniques
4. All of the above

4) It is only the beekeeper's responsibility to keep the bees safe from pesticide applications.

1. True
2. False

Answers

1: 2 2: 1 3: 4 4: 2

Interior Plantscapes

(Some material in this chapter adapted from Michigan State University Extension Bulletin E-2308 Interiorscape Pest Management. A Training Manual for Commercial Pesticide Applicators. Julie Stachecki, Editor.)

Interior plantscapes may be more susceptible to certain diseases and pests due to environmental conditions and fewer natural controls. Diseases on indoor landscape plants are typically a result of adverse environmental conditions combined with the presence of pathogens (fungi, water molds, bacteria, viruses or nematodes). Diseases are uncommon in the indoor landscape setting if it was established using disease-free plant material. It is critical to identify the causal agent or the type of disease that arises in the interior landscape in order to take appropriate corrective measures.



Most pest problems on interiorscape plants originate because the plants were not grown in an indoor environment. Problems typically originate where the plants were originally grown, such as fields, shade houses, or greenhouses. Once introduced to the interiorscape, many insects and mites will thrive, and spread to other plants. The key to preventing insects and mite infestations is to make sure the plants are pest-free before permanent installation. There is much greater flexibility in control procedures (chemicals, application methods, etc.) in a greenhouse or even an acclimatization room than in the interiorscape. Quarantine purchased plants in an isolated

room and carefully inspect and monitor them for insects and mites. If present, use an insecticide or miticide and re-inspect them before moving them to the interiorscape. Avoiding pest problems is often much easier the controlling an established problem.

An amazing variety of insects feed on flowering and foliage plants. The routine use of insecticides usually eliminates predaceous insects and mites. However, pests remaining after treatment sometimes tolerate commonly used insecticides. To stay in business, most commercial flower and foliage plant growers must become fairly sophisticated in using various types of pest management practices, insecticide formulations, application equipment, and in rotating insecticides from one chemical group to another.

The importance of early detection and diagnosis of the problem cannot be overemphasized. This is the key to controlling nearly all pests before significant plant injury or control expenses occur and while pest populations are low. Pesticides and natural enemies will control most insects and mites, if correct procedures are followed.

Commercial indoor landscape accounts are perhaps the most difficult areas in which to attempt pest control. Interior plantscapes extend from public conservatories to extensive plantings in homes, hotels, office buildings, restaurants, shopping malls, hospitals, schools, and other environmentally sensitive areas. The use of insecticides in these areas is often greatly restricted because of the sensitivity of the surroundings. In addition, few chemicals are cleared for ornamental plant use in public areas, and public prejudice against pesticide odors can prevent application of pesticides in many situations.

Most plant replacements result from a combination of poor environmental conditions and poor cultural practices. Symptoms of abiotic disorders, resulting from poor environmental conditions and cultural practices, are extremely variable and are often misdiagnosed. Pesticide applications will not correct an abiotic disorder. It is vital that plant technicians understand the impact of their cultural practices on the plants they maintain.

Prevention

Always inspect new plants for potential pest and disease problems before they are introduced into the interiorscape. Avoid using plants that appear unhealthy or damaged. These problems should be corrected before their use in an interiorscape. Become familiar with the potential pest and disease problems with the types of plants you manage. Highly susceptible plants can serve as indicator plants to alert you to developing problems.

Sanitation

Sanitation is an important component of an integrated approach to pest and disease control. This term refers to the cleanliness approach to excluding and eradicating diseases and pests to eliminate or reduce spread. Exclusion is the first step of a good sanitation program. Avoid introduction of infected or infested material. Once diseases or pests are confirmed in an interiorscape, sanitation methods can reduce their populations and limit spread to healthy plant material. Affected plant tissue should be removed promptly. Heavily infested or diseased plants need to be removed from the interiorscape to reduce spread to other healthier plants. Plant debris should be promptly collected and disposed of. Tools used to prune, replant and care for plants need to be regularly cleaned and sanitized.



Biological Control

Interest in biological control in indoor landscapes has expanded in recent years because of restrictions placed on interiorscape pesticide applications, pesticide costs, poor control with pesticide products, phytotoxicity, need

to repeat applications, and potential human health hazards. Biological control relies on parasitoids, predators, or pathogens to suppress insect and mite pest populations, as well as some fungal diseases. Biological agents are usually released and become established before pest population increase to damaging levels. For some biocontrol agents, they may need to be released periodically. To be successful at managing pest problems with biological control agents requires a proper identification of pests, knowledge of the biology of the pest species, the biological control agent(s), and a great deal of time and commitment. Biological control programs do not look after themselves and need to be maintained properly to be successful.

Host Plant Resistance

Selecting plant species or varieties that are less susceptible to recurring pests and diseases can help to reduce their impact and reliance on pesticides. Proper horticultural care of the plants will also minimize plant stress and help plants better maintain their own defenses. Be careful to avoid over- or under-watering of plants. Intiorscapes are inherently stressful on plants, so try to minimize those stresses on plants. Plants may need to be swapped out so that they can recover from long periods in an interiorscape.

Practice Questions

- 5) Pest problems with interiorscape plants originate from where the plants were originally grown.
1. True
 2. False
- 6) Most pesticides for field use are available for ornamental plant use in public areas.
1. True
 2. False
- 7) Diseases are common with indoor landscape setting if it was established using disease-free plant material.
1. True
 2. False

Answers

1: 1 2: 2 3: 2

Educational programs of Kentucky Cooperative Extension serve all people regardless of economic or social status and will not discriminate on the basis of race, color, ethnic origin, national origin, creed, religion, political belief, sex, sexual orientation, gender identity, gender expression, pregnancy, marital status, genetic information, age, veteran status, or physical or mental disability. Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Nancy M. Cox, Director, Land Grant Programs, University of Kentucky College of Agriculture, Food and Environment, Lexington, and Kentucky State University, Frankfort. Copyright © 2022 for materials developed by University of Kentucky Cooperative Extension. This publication may be reproduced in portions or its entirety for educational or nonprofit purposes only. Permitted users shall give credit to the author(s) and include this copyright notice.